

POISON

**KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

ERASER 750

INSECTICIDE

**ACTIVE CONSTITUENT: 750 g/kg ACEPHATE
(an anticholinesterase compound)**

GROUP 1B INSECTICIDE

For the control of certain insect pests on Bananas,
Crucifers, Macadamias, Ornamentals, Potatoes, Tomatoes
and Tobacco as per the Directions for Use table.



Sinon Australia Pty Limited ACN 102 741 024
1.03/33 Lexington Drive, Bella Vista NSW 2153
TEL: 02 8883 0239

**CONTENTS
5 KG**

APVMA APPROVAL NO: 59572/0306

DIRECTIONS FOR USE

CROP	PEST	STATES	RATE		WHP	CRITICAL COMMENTS
			per 100L water	per ha		
Bananas (Bell Injection and bunch spray only)	Banana Scab moth (<i>Nacoleia octasema</i>) Banana flower thrips (<i>Thrips hawaiiensis</i>)	Qld, NSW, WA, NT only	168g (17g/10L)	-	-	Apply 30mL of prepared solution to each emerging banana bell. Inject the required volume directly into the upright bell as it emerges from the throat of the banana plant.
	Banana rust thrips (<i>Chaetanapho-thrips signipennis</i>)					Apply to the point of run-off, up to 50mL of prepared spray solution to each bunch, ensuring thorough, even spray coverage. Application should be made no later than 2 weeks after bunch emergence.
Brussels Sprouts	Heliothis (<i>Helicoverpa</i> spp.) including corn earworm and native budworm	Tas, NT only	130g	1.3kg	3 days (broccoli 14 days)	Spray when infestation occurs and then at 10 to 14 day intervals or as necessary. Add wetting agent to solution at rate of 10ml per 100 litres.
	Cabbage moth, cabbage white butterfly	NSW, ACT, Vic, SA, Tas, WA, NT only	130g	1.3kg		
	Corn earworm, native budworm	Qld, WA only	100g	1kg		
	Cabbage aphid	Qld, WA, NT only	100g	1kg		
Cabbages, cauliflowers, broccoli	Cabbage moth, cabbage white butterfly	NSW, ACT, Vic, SA, Tas, WA, NT only	130g	1.3kg		
	Cabbage aphid, corn earworm, native budworm	Qld, WA, NT only	100g	1kg		
Macadamias	Flower eating caterpillar, fruitspotting bug, macadamia leaf miner, banana spotting bug, flower thrips	Qld, NT only	66g	660g	-	Apply at early flowering if pests are evident. One to three sprays may be required depending on the time and severity of the infestation. Spray when bees are least active.
	Monolepta beetle	Qld, WA, NT only	100g	1kg		Apply when activity first observed and repeat at 14 to 21 day intervals or as necessary.
	Fruitspotting bug, macadamia flower caterpillar, macadamia leaf miner, Macadamia nutborer	NSW, WA only				
Ornamentals	Aphids	All States	130g	1.3kg	-	Spray when infestation occurs and then at 14 day intervals. DO NOT apply to carnations later than 6 days before picking.
	Corn earworm, Native budworm	Vic, SA, Tas, WA only				
Potatoes	Potato moth (tobacco leafminer)	All States	66g	660g	3 days	Spray every 14 to 21 days or as necessary.
Tobacco	Tobacco budworm	NSW, ACT only	130g	1.3kg	14 days	Spray at 7 to 10 day intervals, when the pest is present.
	Cluster caterpillar, Heliothis (<i>Helicoverpa</i> spp.) Tobacco looper, Tobacco stemborer, White flies	Qld, WA only				Spray as indicated by crop checking (usually a minimum of 7 days is required). Use high volume application for good coverage ie 130 to 400L/ha depending on the crop stage, paying particular attention to bud heart.
	Heliothis (<i>Helicoverpa</i> spp.)	Vic only	100g	1kg		
Tomatoes	Aphids	NSW, ACT, Vic, SA, WA, Tas only	100g	1kg	3 days	Spray when infestation occurs and then at 7 to 14 day intervals or as necessary (ground application only)
	Potato moth, Heliothis (<i>Helicoverpa</i> spp.)		130g	1.3kg		
	Potato moth, Heliothis (<i>Helicoverpa</i> spp.), aphids	Qld, NT only	100g	1kg		Apply when infestation occurs then at 7 day intervals or as necessary.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD:

BRUSSELS SPROUT, CABBAGE, CAULIFLOWERS, POTATOES AND TOMATOES – DO NOT HARVEST FOR 3 DAYS AFTER APPLICATION.

BROCCOLI, TOBACCO – DO NOT HARVEST FOR 14 DAYS AFTER APPLICATION.

BANANAS, MACADAMIAS, ORNAMENTALS – NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

MIXING

Add the required amount of Eraser 750 Insecticide to the full volume of water in the spray tank with agitator in operation. Ensure the powder is completely dispersed before spraying.

NOTE: A varying degree of consolidation of the contents of this container may occur during transport but the net weight of the contents is not changed and is as stated on the label.

APPLICATION METHOD

By ground:

Crops should be sprayed with sufficient volume to ensure adequate coverage of plants. For high volume spraying, apply in 500 to 1000 litres water per hectare.

By air:

Apply in 10 to 45 litres of water per hectare.

INSECTICIDE RESISTANCE WARNING

GROUP 1B INSECTICIDE

For insecticide resistance management Eraser 750 Insecticide is a Group 1B insecticide.

Some naturally occurring insect biotypes resistant to Eraser 750 Insecticide and other Group 1B Insecticides may exist through normal genetic variability in any insect population. The resistant individuals can eventually dominate the insect population if these insecticides are used repeatedly. The effectiveness of Eraser 750 Insecticide on resistant individuals could be significantly reduced. Since the occurrence of resistant individuals is difficult to detect prior to use, Sinon Australia Pty Limited accepts no liability for any losses that may result from the failure of Eraser 750 Insecticide to control resistant insects.

Eraser 750 Insecticide may be subject to specific resistance management strategies. For further information contact your local supplier, Sinon representative or local agricultural department agronomist.

COMPATIBILITY

Eraser 750 Insecticide may be mixed in the spray vat with any of the following products: Copper Oxychloride, Zineb. A mixture of Eraser 750 Insecticide with other products may be ineffective or may cause serious damage and hence is NOT recommended.

PROTECTION OF CROPS, NATIVE AND OTHER NON-TARGET PLANTS

Eraser 750 Insecticide MAY CAUSE FOLIAGE DAMAGE TO CERTAIN SHRUBS AND TREES i.e. Crab Apple, Cotton wood, Lombardy poplar and American elm. DO NOT spray more than once every 28 days on carnations and chrysanthemums. Certain chrysanthemum varieties ie Showoff, Iceberg, Albatross, Bonnie Jean and Statesman have, on occasions, been damaged by products containing acephate.

PROTECTION OF LIVESTOCK, WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

Dangerous to bees. DO NOT spray any plants in flower while bees are foraging. Avoid spray drift from target area. DO NOT contaminate streams, rivers or waterways with this product or the used containers.

STORAGE AND DISPOSAL

Store in the original container, tightly closed in a safe well-ventilated area, as cool as possible. Do not store for prolonged periods in direct sunlight. Single rinse before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site. Puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500mm below the surface in a disposal pit specifically marked and set up

for this purpose clear of waterways, desirable vegetation and tree roots, in compliance with relevant Local, State or Territory government regulations. Do not burn empty containers and product.

SAFETY DIRECTIONS

Product and spray are poisonous if absorbed by skin contact, inhaled or swallowed. Repeated minor exposure may have a cumulative poisoning effect. Avoid contact with eyes and skin. Do not inhale dust or spray mist.

When preparing the spray and using the prepared spray, wear cotton overalls buttoned to the neck and wrist, a washable hat, elbow-length PVC gloves and face shield. After use and before eating drinking or smoking, wash hands, arms, and face thoroughly with soap and water. After each day's use, wash gloves, face shield and contaminated clothing.

FIRST AID

If swallowed, splashed on skin or in eyes, or inhaled, contact a doctor or Poisons Information Centre (Phone Australia 13 11 26) or a doctor at once. Remove any contaminated clothing and wash skin thoroughly. If swallowed, activated charcoal may be advised. Give atropine if instructed.

MATERIAL SAFETY DATA SHEET

For further information refer to the Material Safety Data Sheet (MSDS) which is available from the supplier.

NOTICE TO BUYER

Sinon Australia Pty Limited will not be held liable for any loss, injury or damage, indirect or consequential, arising from the sale, supply, use or application of this product. The product is not to be used for any purpose or in any way contrary to label instructions.

**ORGANOPHOSPHORUS PESTICIDE,
SOLID, TOXIC, N.O.S.
(contains Acephate)**

UN No.2783

**IN A TRANSPORT EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**



Date of Manufacture:

Batch No:

ERASERKG-0005-0813
APVMA Approval No: 59572/0306

